



**VYALEV™** COMPLETE



## Traveling with your delivery system

Find out what to expect  
and how to prepare

Please see Use and Important Safety Information on page 8.  
Please visit [https://www.rxabbvie.com/pdf/vyalev\\_pi.pdf](https://www.rxabbvie.com/pdf/vyalev_pi.pdf) for  
full Prescribing Information.



**VYALEV™**

**foscarbidopa/foslevodopa**

Injection for subcutaneous use  
12 mg/240 mg per mL



**Talk to your Nurse Ambassador\* for tips on traveling with sharps and other used supplies.**


\*Nurse Ambassadors are provided by AbbVie and do not work under the direction of your health care professional (HCP) or give medical advice. They are trained to direct patients to their HCP for treatment-related advice, including further referrals.

Please see Use and Important Safety Information on [page 8](#).  
Please visit [https://www.rxabbvie.com/pdf/vyalev\\_pi.pdf](https://www.rxabbvie.com/pdf/vyalev_pi.pdf) for  
full Prescribing Information.



## Packing Your Supplies

You'll be wearing your pump in its carrying case when you're traveling. But whether you're taking a long vacation or going away for the weekend, you'll need to bring your other supplies, too. **Be sure to pack enough to last the whole time:**

- ☐ Syringes
- ☐ Infusion sets
- ☐ Medication vials
- ☐ Vial adapters
- ☐ Alcohol pads
- ☐ Fully charged spare battery
- ☐ Battery charging station
- ☐ Your TSA Notification & Medical Alert cards
- ☐ Your prescribed oral carbidopa-levodopa immediate-release tablets, if you're unable to use your delivery system
-  Consider bringing your How-to Guide



Your medication vials can be stored at room temperature for up to **28 days**. Once VYALEV has been stored at room temperature, do not return the product to the refrigerator. Discard after 28 days. Do not freeze the medication. If you're traveling within the US for more than **28 days**, ask your Specialty Pharmacy<sup>†</sup> about shipping your medication to your destination.

<sup>†</sup>A Specialty Pharmacy is a type of pharmacy that handles medication for conditions like advanced Parkinson's. These medications often need special storage and processing.

Please see Use and Important Safety Information on [page 8](#).  
Please visit [https://www.rxabbvie.com/pdf/vyalev\\_pi.pdf](https://www.rxabbvie.com/pdf/vyalev_pi.pdf) for  
full Prescribing Information.



## Flying & TSA screening

Your delivery system can work on your flight. But your pump and some of your supplies can't go through the screening machines at security. You'll need to request a different screening process so **you don't have to disconnect from your pump.**



### **CAN** go through screening machines

- Unopened supplies:
  - Infusion sets
  - Vial adapters
  - Medication vials
  - Syringes
  - Alcohol pads
- Prescribed carbidopa-levodopa tablets

Pack these supplies in your carry-on bag. They should not be packed in checked luggage, which could get lost, delayed or exposed to damaging temperatures.



### **CANNOT** go through screening machines

- Pump, in its carrying case
- Pump batteries
- Charging station

You'll need to remove your charging station and spare battery from your carry-on bag before it goes through the machines. Consider packing them separately from your other supplies.



## Flying & TSA screening *cont'd*



### Call the Transportation Security Administration (TSA) 72 hours before your flight

Tell them about your treatment and let them know you'll need an alternative screening process. You can reach them at **855-787-2227**.

## Things to do on the day of your flight

- ☐ **Replace your infusion supplies before heading out.** So you might not have to change them at the airport or on the plane
- ☐ **Remind TSA personnel that you'll need an alternative screening process.** Show them your TSA Notification Card and your Medical Alert Card
- ☐ **Arrive at the airport early.** You'll need plenty of time to go through security
- ☐ **Ask to board the plane early so you can keep your carry-on bags with you.** Your supplies should not go in checked luggage

Please see Use and Important Safety Information on [page 8](#).  
Please visit [https://www.rxabbvie.com/pdf/vyalev\\_pi.pdf](https://www.rxabbvie.com/pdf/vyalev_pi.pdf) for  
full Prescribing Information.



# Your travel cards

## Don't forget to bring these cards

Print out this page and cut out the front and back of the TSA Notification and Medical Alert cards. You can keep them in your wallet, pump carrying case or somewhere easy to access.

### TSA Notification Card

NAME: \_\_\_\_\_

**I have a health condition that requires a medical device. This may affect my screening.**

My treatment requires I carry an ambulatory pump that delivers a subcutaneous infusion into my body 24 hours a day through a cannula attached to my abdomen.

### I understand:

- This card does not exempt me from screening.
- Alternative screening procedures are available and can be done in private.
- I may ask to speak to a supervisory TSA officer.

For additional questions, call TSA Cares at 1-855-787-2227.

© 2024 AbbVie. All rights reserved.  
US-LEVCAR-220080 October 2024

Your **TSA Notification Card** will let airport security know you can't go through the screening machines.



Cut out the front and back along the dotted lines.

### Medical Alert Card

My Name Is: \_\_\_\_\_

- I am receiving therapy called VYALEV (foscarbidopa/foslevodopa).
- VYALEV is delivered with a pump.
- There is no immediate risk to me if you disconnect from the infusion site or remove the cannula from my abdomen.

### In case of emergency, please contact:

Emergency Contact: \_\_\_\_\_

Primary Care Doctor: \_\_\_\_\_

VYALEV Prescriber: \_\_\_\_\_

VYALEV Complete  
24/7 Hotline: \_\_\_\_\_

1-866-489-2538

Your **Medical Alert Card** will let people know about your treatment and who to contact in case of an emergency.



Cut out the front and back along the dotted lines.

Please see Use and Important Safety Information on **page 8**.  
Please visit [https://www.rxabbvie.com/pdf/vyalev\\_pi.pdf](https://www.rxabbvie.com/pdf/vyalev_pi.pdf) for full Prescribing Information.





## Have any questions?

Reach out to your Ambassador for more tips and resources on traveling. If they're not available, call the **VYALEV Complete 24/7 Hotline<sup>‡</sup> at 1-866-489-2538.**

<sup>‡</sup>The 24/7 Hotline is staffed by nurses that are specifically trained on VYALEV and can be reached 24 hours per day, 7 days a week. After business hours, you may be asked to leave a message for a return call within 30 minutes.

Please see Use and Important Safety Information on **page 8**.  
Please visit [https://www.rxabbvie.com/pdf/vyalev\\_pi.pdf](https://www.rxabbvie.com/pdf/vyalev_pi.pdf) for  
full Prescribing Information.



## Use

VYALEV is a prescription medicine used for treatment of advanced Parkinson's disease in adults. VYALEV contains two medicines, foscarnidopa and foslevodopa.

## Important Safety Information

**What is the most important safety information I should know about VYALEV™ (foscarnidopa/foslevodopa)?**

Do not take VYALEV if you currently take or have recently taken (within the last 14 days) a medication for depression called a nonselective monoamine oxidase (MAO) inhibitor. Ask your healthcare provider or pharmacist if you are not sure if you take an MAO inhibitor.

**Tell your healthcare provider about all your medical conditions and the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. When used together, VYALEV and certain other medicines, including medications for high blood pressure, MAO inhibitors, antipsychotics, metoclopramide, and isoniazid, may affect each other and cause serious side effects.**

**VYALEV may cause other serious side effects. Talk to your healthcare provider before starting VYALEV and while on VYALEV if you have had or have any of the following:**

- **Falling asleep without warning during normal daily activities.** VYALEV may cause you to fall asleep while you are doing daily activities, such as driving, which may result in an accident. This can happen as late as 1 year after you start VYALEV. Do not drive or operate machinery until you know how VYALEV affects you. Tell your healthcare provider if you take medicines that can make you sleepy, such as sleep medicines, antidepressants, or antipsychotics.
- **Seeing, hearing, or feeling things that are not real (hallucinations).** This is a common and sometimes serious side effect.
- **Unusual urges.** Some people taking medicines for Parkinson's disease, including VYALEV, have reported problems, such as gambling, compulsive eating, compulsive shopping, and increased sex drive.

- **Infusion Site Reactions and Infections.** Some people using VYALEV have had reactions and infections at the infusion site. **This is a common and sometimes serious side effect.** Remove your cannula and **call your healthcare provider if you have any of the following symptoms of an infection:** local spreading of redness, pain, swelling, warmth, change in color when pressing area, or fever. Take oral carbidopa/levodopa tablets until you are able to resume VYALEV.
- **Uncontrolled sudden movements (dyskinesia).** **This is a common and sometimes serious side effect.** If you have new dyskinesia or your dyskinesia gets worse, tell your healthcare provider. This may be a sign that your dose of VYALEV or other Parkinson's medicines may need to be adjusted.
- **Heart attack or other heart problems.** Tell your healthcare provider if you have had increased blood pressure, a fast or irregular heartbeat, or chest pain.
- Worsening of the **increased pressure in your eyes** (glaucoma). The pressure in your eyes should be checked after starting VYALEV.

**Do not stop using VYALEV or change your dose unless you are told to do so by your healthcare provider. Tell your healthcare provider if you develop withdrawal symptoms, such as fever, confusion, or severe muscle stiffness.**

These are not all the possible side effects of VYALEV. For more information, ask your healthcare provider or pharmacist.

VYALEV (foscarnidopa and foslevodopa) injection for subcutaneous use is available in a 120 mg foscarnidopa and 2,400 mg foslevodopa per 10 mL (12 mg foscarnidopa and 240 mg foslevodopa per mL) solution.

**Please see the full Prescribing Information, including Medication Guide, for additional information about VYALEV. Talk to your healthcare provider if you have questions.**

**You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088.**

**If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit [AbbVie.com/PatientAccessSupport](http://AbbVie.com/PatientAccessSupport) to learn more.**

Please visit [https://www.rxabbvie.com/pdf/vyalev\\_pi.pdf](https://www.rxabbvie.com/pdf/vyalev_pi.pdf) for full Prescribing Information.

abbvie