

Use this mat to get ready for your VYALEV treatment

Let's lay it all out together

Refer to the Patient Instructions for Use materials that came with your drug vials, pump and other supplies from your Specialty Pharmacy.*

Follow the Plan-it-Out steps:

- 1 Wash your hands thoroughly with soap and water for at least 20 seconds. Scrub the back of your hands, between your fingers and under your nails. Dry with a new paper towel.
- 2 Grab your medication vial. If it was in the fridge, let it sit at room temperature for ⌚ 30 minutes before using it.
 - If refrigerated, do not warm VYALEV (in solution vial or syringe) in any way other than letting it warm at room temperature. For example, do not warm in a microwave or in hot water.
- 3 Clean a large, flat workspace.
- 4 Gather your supplies:
 - ⚠ Check for packaging damage and expiration dates. Do not use any damaged or expired supplies
 - 💊 Inspect the medication vial. Do not use it if there are particles, cloudiness or flakes
 - 🔋 Insert a charged battery into your pumpKeep the supplies in their packaging until you're ready to use them.
- 5 Wash and dry your hands again before handling your supplies.

*A Specialty Pharmacy is a type of pharmacy that handles medications that often need special storage and processing.


Please see Use and Important Safety Information on [page 4](#), and full [Prescribing Information](#).



VYALEV™
foscarbidopa/foslevodopa
Injection for subcutaneous use
12 mg/240 mg per mL


Paper Towels

You'll need a few new paper towels when priming the infusion set.



Vial Adapter

Allows the syringe to remove the medication from the vial.



Infusion Set

Tubing


Delivers medication from the syringe to the infusion site.

Cannula Insertion Device

Inserts the cannula into the skin of your belly.



PUT A FULLY CHARGED BATTERY IN YOUR PUMP BEFORE YOU START



Pump

Delivers the medication. Comes with 2 batteries. Charge one battery while the other is in use.


Alcohol Pads

Used to swab the top of the medication vial before use and to swab your skin before inserting the cannula.



Medication Vial

Check the solution. It shouldn't be cloudy or have particles or flakes.



Syringe

Used to draw out the medication from the vial. Then placed in the pump.



Helping you navigate your treatment journey

We're here to support you every step of the way

Check out these resources:



- **Help from your Nurse Ambassador***: Give them a call. If they're not available, call the VYALEV Complete 24/7 Hotline† at **1-866-489-2538**



- Visit www.VYALEV.com

Refer to the Patient Instructions for Use materials that came with your drug vials, pump and other supplies from your Specialty Pharmacy.‡

*Nurse Ambassadors are provided by AbbVie and do not work under the direction of your health care professional (HCP) or give medical advice. They are trained to direct patients to their HCP for treatment-related advice, including further referrals.

†The 24/7 Hotline is staffed by nurses that are specifically trained on VYALEV and can be reached 24 hours per day, 7 days a week. After business hours, you may be asked to leave a message for a return call within 30 minutes.

‡A Specialty Pharmacy is a type of pharmacy that handles medications that often need special storage and processing.

Please see Use and Important Safety Information on [page 4](#), and full [Prescribing Information](#).



VYALEV™
foscarbidopa/foslevodopa
Injection for subcutaneous use

Use

VYALEV is a prescription medicine used for treatment of advanced Parkinson's disease in adults. VYALEV contains two medicines, foscarnidopa and foslevodopa.

Important Safety Information

What is the most important safety information I should know about VYALEV™ (foscarnidopa/foslevodopa)?

Do not take VYALEV if you currently take or have recently taken (within the last 14 days) a medication for depression called a nonselective monoamine oxidase (MAO) inhibitor. Ask your healthcare provider or pharmacist if you are not sure if you take an MAO inhibitor.

Tell your healthcare provider about all your medical conditions and the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. When used together, VYALEV and certain other medicines, including medications for high blood pressure, MAO inhibitors, antipsychotics, metoclopramide, and isoniazid, may affect each other and cause serious side effects.

VYALEV may cause other serious side effects. Talk to your healthcare provider before starting VYALEV and while on VYALEV if you have had or have any of the following:

- **Falling asleep without warning during normal daily activities.** VYALEV may cause you to fall asleep while you are doing daily activities, such as driving, which may result in an accident. This can happen as late as 1 year after you start VYALEV. Do not drive or operate machinery until you know how VYALEV affects you. Tell your healthcare provider if you take medicines that can make you sleepy, such as sleep medicines, antidepressants, or antipsychotics.
- **Seeing, hearing, or feeling things that are not real (hallucinations). This is a common and sometimes serious side effect.**
- **Unusual urges.** Some people taking medicines for Parkinson's disease, including VYALEV, have reported problems, such as gambling, compulsive eating, compulsive shopping, and increased sex drive.
- **Infusion Site Reactions and Infections.** Some people using VYALEV have had reactions and infections at the infusion site. **This is a common**

and sometimes serious side effect. Remove your cannula and **call your healthcare provider if you have any of the following symptoms of an infection:** local spreading of redness, pain, swelling, warmth, change in color when pressing area, or fever. Take oral carbidopa/levodopa tablets until you are able to resume VYALEV.

- **Uncontrolled sudden movements (dyskinesia). This is a common and sometimes serious side effect.** If you have new dyskinesia or your dyskinesia gets worse, tell your healthcare provider. This may be a sign that your dose of VYALEV or other Parkinson's medicines may need to be adjusted.
- **Heart attack or other heart problems.** Tell your healthcare provider if you have had increased blood pressure, a fast or irregular heartbeat, or chest pain.
- Worsening of the **increased pressure in your eyes (glaucoma).** The pressure in your eyes should be checked after starting VYALEV.

Do not stop using VYALEV or change your dose unless you are told to do so by your healthcare provider. Tell your healthcare provider if you develop withdrawal symptoms, such as fever, confusion, or severe muscle stiffness.

These are not all the possible side effects of VYALEV. For more information, ask your healthcare provider or pharmacist.

VYALEV (foscarnidopa and foslevodopa) injection for subcutaneous use is available in a 120 mg foscarnidopa and 2,400 mg foslevodopa per 10 mL (12 mg foscarnidopa and 240 mg foslevodopa per mL) solution.

Please see the full Prescribing Information, including Medication Guide, for additional information about VYALEV. Talk to your healthcare provider if you have questions.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

If you are having difficulty paying for your medicine, AbbVie may be able to help.

Visit AbbVie.com/PatientAccessSupport to learn more.

abbvie



VYALEV™
foscarnidopa/foslevodopa
Injection for subcutaneous use